



Veronese transform, and Castelnuovo-Mumford regularity of modules

Marcel Morales, Dung Nguyen Thi

► To cite this version:

Marcel Morales, Dung Nguyen Thi. Veronese transform, and Castelnuovo-Mumford regularity of modules. Turkish Journal of Mathematics, 2016, 40 (4), pp.838-849. hal-00963895

HAL Id: hal-00963895

<https://hal.science/hal-00963895>

Submitted on 22 Mar 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Veronese transform, and Castelnuovo-Mumford regularity of modules

MARCEL MORALES

Université de Grenoble I, Institut Fourier, UMR 5582, B.P.74,
38402 Saint-Martin D'Hères Cedex,
and IUFM de Lyon, 5 rue Anselme,
69317 Lyon Cedex (FRANCE)

NGUYEN THI DUNG

Thai Nguyen University of Agriculture and Forestry,
Thai Nguyen,
Vietnam,

Abstract ¹

Veronese rings, Segre embeddings or more generally Segre-Veronese embeddings are very important rings in Algebraic Geometry. In this paper we present an original, elementary way to compute the Hilbert-Poincaré series of these rings, as a consequence we compute their Castelnuovo-Mumford regularity, and also the leading term of the h -vector. Moreover, we can compute the Castelnuovo-Mumford regularity of the n -Veronese Module of any finitely generated Cohen-Macaulay graded module.

1 Introduction

Veronese rings, Segre embeddings or more generally Segre-Veronese embeddings are very important rings in Algebraic Geometry. It is well known that these rings are arithmetically Cohen-Macaulay, hence their Hilbert-Poincaré series can be written $P_R(t) = \frac{Q_R(t)}{(1-t)^{\dim R}}$, where $Q_R(t)$ is a polynomial on t with $Q_R(1) \neq 0$ having positive integer coefficients, the sequence of the coefficients of $Q_R(t)$ is also called the h -vector of R . The degree of $Q_R(t)$ is the Castelnuovo-Mumford regularity (c.f.[7][Chapter 4]), and the leading term of $Q_R(t)$ is the highest graded Betti number of R . By using very original and elementary methods we are able to compute the leading term of $Q_R(t)$. Our results allows to compute the Castelnuovo-Mumford regularity of the n Veronese Module of any finitely generated Cohen-Macaulay graded module, and the rings called of Veronese type.

¹ Partially supported by VIASM, Hanoi, Vietnam.

MSC 2000: Primary: 13D02, Secondary 14F05.

Key words and phrases: Castelnuovo-Mumford regularity, Veronese ring, Segre ring, Hilbert Series.

Note that this result can be proved easily by using local cohomology, but our purpose is to give a very elementary proof.

Our main results improves partially [1] and [5].

Theorem. *Let consider the Segre-Veronese ring $R_{\underline{b}, \underline{n}}$, $\dim R_{\underline{b}, \underline{n}} = b_1 + \dots + b_m + 1$. Let $P_{R_{\underline{b}, \underline{n}}}(t) = \frac{Q_{R_{\underline{b}, \underline{n}}}}{(1-t)^{\dim R_{\underline{b}, \underline{n}}}}$ be the Hilbert-Poincaré series of $R_{\underline{b}, \underline{n}}$, with $Q_{R_{\underline{b}, \underline{n}}} = h_0 + h_1 t + \dots + h_{r_{\underline{b}, \underline{n}}} t^{r_{\underline{b}, \underline{n}}}$, where $r_{\underline{b}, \underline{n}}$ is the Castelnuovo-Mumford regularity of $R_{\underline{b}, \underline{n}}$. We set $\alpha_{\underline{b}, \underline{n}} = \dim R_{\underline{b}, \underline{n}} - r_{\underline{b}, \underline{n}}$. After a permutation of b_1, \dots, b_m , we can assume that, for all $i = 1, \dots, m$, $\lceil \frac{b_1+1}{n_1} \rceil > \frac{b_i}{n_i}$. Then*

$$\alpha_{\underline{b}, \underline{n}} = \lceil \frac{b_1 + 1}{n_1} \rceil, \quad ,$$

and the highest Betti number of $R_{\underline{b}, \underline{n}}$ is

$$\beta_{r_{\underline{b}, \underline{n}}} = h_{r_{\underline{b}, \underline{n}}} = \binom{n_1 \alpha_{\underline{b}, \underline{n}} - 1}{b_1} \dots \binom{n_m \alpha_{\underline{b}, \underline{n}} - 1}{b_m}$$

Theorem. *Fix integers $d, n \in \mathbb{N}^*, \tau \in \mathbb{Z}$. Let $(a_l)_{l \in \mathbb{Z}}$ be a sequence of complex numbers, such that $a_l = 0$ for $l \ll 0$, set :*

$$f(t) = \sum_{l \in \mathbb{Z}} a_l t^l, \quad f^{<n, \tau>}(t) = \sum_{l \in \mathbb{Z}} a_{nl + \tau} t^l.$$

If $f(t) = \frac{h(t)}{(1-t)^d}$ with $h(t) \in \mathbb{C}[t, t^{-1}]$ then $f^{<n, \tau>}(t) = \frac{h^{<n>}(t)}{(1-t)^d}$ with $h^{<n>}(t) \in \mathbb{C}[t, t^{-1}]$ such that:

- $\deg h^{<n, \tau>}(t) \leq d - \lceil \frac{d - \deg h(t) + \tau}{n} \rceil,$
- *If all the coefficients of $h(t)$ are positive real numbers then*
 $\deg h^{<n, \tau>}(t) = d - \lceil \frac{d - \deg h(t) + \tau}{n} \rceil,$
- *If $\deg h(t) = d$ then $\deg h^{<n>}(t) = d$.*

2 Preliminaries on toric rings and Hilbert-Poincaré series

Let $S = K[x_0, \dots, x_s, x_0^{-1}, \dots, x_s^{-1}]$ be a Laurent polynomial ring over a field K on a finite set of variables. For any finite set \mathcal{M} of monomials in S , let $K[\mathcal{M}] \subset S$ be the subring of S generated by the set

\mathcal{M} . It is the toric ring defined by the semigroup generated by \mathcal{M} . In what follows we consider the special case where $S = K[x_0, \dots, x_s]$ is a polynomial ring over the field K and all the monomials in \mathcal{M} are of the same degree.

Example 2.1. Let $S = K[x_0, \dots, x_b] \oplus_{l \in \mathbb{N}} S_l$, and $\mathcal{M} = \{x_0^{\alpha_0} \dots x_b^{\alpha_b} \mid \alpha_0 + \dots + \alpha_b = n\}$. So that

$$R_{b,n} = K[\mathcal{M}] = \oplus_{l \in \mathbb{N}} S_{nl}.$$

This toric ring is known as the n -Veronese embedding of S .

Example 2.2. More generally, let X_1, \dots, X_m , m sets of independent disjoint variables, with $\text{Card}(X_i) = b_i + 1$. Let $S_i = K[X_i]$ for $i = 1, \dots, m$, $S = K[X_1 \cup X_2 \cup X_m]$, and $\mathcal{M} = \{x_1 x_2 \dots x_m \mid x_i \in X_i\}$. So that

$$R_{b_1, \dots, b_m} = K[\mathcal{M}] = \oplus_{l \in \mathbb{N}} (S_1)_l \otimes \dots \otimes (S_m)_l.$$

This toric ring is known as the Segre embedding of the m polynomial rings S_1, \dots, S_m .

Example 2.3. Let X_1, \dots, X_m , sets of independent disjoint variables such that $X_i = \{x_{i,0}, \dots, x_{i,b_i}\}$, $S_i = K[X_i]$ for $i = 1, \dots, m$, and $n_1, \dots, n_m \in \mathbb{N}$, Let $S = K[X_1 \cup X_2 \cup X_m]$, and

$$\mathcal{M} = \{\underline{x}_1^{\alpha_1} \dots \underline{x}_m^{\alpha_m} \mid |\alpha_i| = n_i\},$$

where $\alpha_i = (\alpha_{i,0}, \dots, \alpha_{i,b_i})$, $\underline{x}_i^{\alpha_i} = x_{i,0}^{\alpha_{i,0}} \dots x_{i,b_i}^{\alpha_{i,b_i}}$, and $|\alpha_i| = \alpha_{i,0} + \dots + \alpha_{i,b_i}$. The Segre-Veronese embedding:

$$R_{\underline{b}, \underline{n}} = K[\mathcal{M}] = \oplus_{l \in \mathbb{N}} (S_1)_{n_1 l} \otimes \dots \otimes (S_m)_{n_m l},$$

where $\underline{b} = (b_1, \dots, b_m)$, $\underline{n} = (n_1, \dots, n_m)$.

Let $S = K[x_0, \dots, x_s]$ be a polynomial ring over the field K , graded by the standard graduation, that is $\deg x_i = 1$, for all i . Let $R := S/I$, where $I \subset S$ is a graded ideal, let $M = \oplus_{l \in \mathbb{Z}} M_l$ be a finitely generated graded R -module, hence M is also a S -module. the Hilbert-function of M is defined by $H_M(l) = \dim_K M_l$, for all $l \in \mathbb{Z}$, and the Hilbert-Poincaré series of M :

$$P_M(t) = \sum_{l \in \mathbb{Z}} H_M(l) t^l.$$

It is well known that

$$P_M(t) = \frac{Q_M(t)}{(1-t)^{\dim M}} ,$$

where $Q_M(t)$ is a Laurent polynomial on t, t^{-1} with $Q_M(1) \neq 0$. Moreover if M is a Cohen-Macaulay S -module, all the coefficients of $Q_M(t)$ are natural integers, and the Castelnuovo-Mumford regularity of M is the degree of $Q_M(t)$. For more details on Hilbert-Poincaré series see [10], [4][Chapter 4], [7][Chapter 4].

Theorem 2.4. (*Hilbert's Theorem*) let $M = \oplus_{l \in \mathbb{Z}} M_l$ be a finitely generated graded S -module. There exists a polynomial with integer coefficients $\Phi_{H_M}(l)$ such that $H_M(l) = \Phi_{H_M}(l)$, for l large enough. Moreover the leading term of $\Phi_{H_M}(l)$ can be written as : $\frac{\deg(M)}{d!} l^d$, where $d+1$ is the dimension of M and $\deg(M)$ the degree or multiplicity of M .

Remark 2.5. The postulation number of the Hilbert function is the biggest integer l such that $H_M(l) \neq \Phi_{H_M}(l)$. It is well known, ([10], [4][Chapter 4]), that the postulation number equals the degree of the rational fraction defining the Poincaré series.

Remark 2.6. We recall that binomial coefficients can be defined in a more general setting than natural numbers, indeed for $k \in \mathbb{N}$, binomial coefficients are polynomial functions in the variable n . More precisely:

(1) If $k = 0$ then let $\binom{n}{0} = 1$, for all $n \in \mathbb{C}$.

(2) If $k > 0$ then let $\binom{n}{k} = \frac{n(n-1)\dots(n-k+1)}{k!}$, for all $n \in \mathbb{C}$.

Note that for all $n \in \mathbb{C}$, $\binom{n}{k} = (-1)^k \binom{k-n-1}{k}$ and if $n \in \mathbb{N}$, $n < k$, then $\binom{n}{k} = 0$.

Example 2.7. Let $S = K[x_0, \dots, x_b]$, be a polynomial ring. Then

$$H_S(l) = \begin{cases} \binom{l+b}{b} & \text{if } l \geq 0 \\ 0 & \text{if } l < 0 \end{cases} , \quad P_S(t) = \frac{1}{(1-t)^{b+1}}.$$

Note that in fact $\forall l \geq -b$, $H_S(l) = \binom{l+b}{b}$ and $0 = H_S(-b-1) \neq \binom{-b-1+b}{b} = (-1)^b$, so the postulation number of S is $-(b+1)$.

Example 2.8. Let $S = K[x_0, \dots, x_b]$, $\mathcal{M} = \{x_0^{\alpha_0} \dots x_b^{\alpha_b} \mid \alpha_0 + \dots + \alpha_b = n\}$, and $R_{b,n} = K[\mathcal{M}]$ the n -Veronese embedding. Then

$$H_{R_{b,n}}(l) = H_S(nl) = \begin{cases} \binom{nl+b}{b} & \text{if } l \geq 0 \\ 0 & \text{if } l < 0 \end{cases}.$$

Note that $\binom{nl+b}{b} = \frac{(nl+1)(nl+2)\dots(nl+b)}{b!}$ is a polynomial on l with leading term $\frac{n^b l^b}{b!}$, so that $\deg(R_{b,n}) = n^b$, $\dim R_{b,n} = b+1$. Note also that $\forall l > -\lceil \frac{b+1}{n} \rceil$, $H_{R_{b,n}}(l) = \binom{nl+b}{b}$ and $0 = H_{R_{b,n}}(-\lceil \frac{b+1}{n} \rceil) \neq \binom{-\lceil \frac{b+1}{n} \rceil + b}{b} = (-1)^b \binom{\lceil \frac{b+1}{n} \rceil - 1}{b}$, so the postulation number of $R_{b,n}$ is $-\lceil \frac{b+1}{n} \rceil$. More generally the postulation number of $R_{b,n}[\tau]$ is $\lceil \frac{b+1+\tau}{n} \rceil$.

3 Veronese of generating series

In a recent paper [2], Brenti and Walker prove that taking the n -Veronese transform of the h polynomial is a linear function, in this section we improve this result giving a elementary proof of the fact that taking the shifted n -Veronese transform of the h polynomial is a linear function on h .

Let recall the following fact:

Theorem 3.1. Let $(a_l)_{l \in \mathbb{Z}}$ be a sequence of complex numbers, such that $a_l = 0$ for $l \ll 0$, set : $f(t) = \sum_{l \in \mathbb{Z}} a_l t^l$, TFAE:

- There exists $h(t) \in \mathbb{C}[t, t^{-1}]$ and a natural integer d such that $f(t) = \frac{h(t)}{(1-t)^d}$.
- There exists $\Phi(t) \in \mathbb{C}[t, t^{-1}]$ of degree $d-1$ with leading coefficient $e_0/(d-1)!$, such that $\Phi(l) = a_l$ for l large enough.

Moreover $h(1) = e_0$.

Definition 3.2. Fix integers $d, n \in \mathbb{N}^*$, $\tau \in \mathbb{Z}$. Let $(a_l)_{l \in \mathbb{Z}}$ be a sequence of complex numbers, such that $a_l = 0$ for $l \ll 0$, set :

$$f(t) = \sum_{l \in \mathbb{Z}} a_l t^l, \quad f^{<n, \tau>}(t) = \sum_{l \in \mathbb{Z}} a_{nl+\tau} t^l.$$

By the Theorem 3.1 if $f(t) = \frac{h(t)}{(1-t)^d}$ with $h(t) \in \mathbb{C}[t, t^{-1}]$ then $f^{<n, \tau>}(t) = \frac{h^{<n, \tau>}(t)}{(1-t)^d}$ with $h^{<n, \tau>}(t) \in \mathbb{C}[t, t^{-1}]$.

Let us introduce some notations. To any non zero polynomial $h(t) = h_\sigma t^\sigma + \dots + h_0 + h_1 t + \dots + h_s t^s \in \mathbb{C}[t, t^{-1}]$ we associate the h -vector $\vec{h} = (\dots, 0, h_\sigma, \dots, h_s, 0, \dots)$, and we set $\deg \vec{h} = \deg h(t)$. For $j \in \mathbb{Z}$, let $\vec{\varepsilon}_j$ be the h -vector of the polynomial t^j . Let denote by $[t^k]h(t)$ the coefficient of t^k in the polynomial $h(t)$. For any $i, j \in \mathbb{Z}$ define $\mathcal{D}_{i,j}$ by

$$\mathcal{D}_{i,j} = [t^{in-j}] \left(\frac{(1-t^n)^d}{(1-t)^d} \right) = [t^{in-j}] ((1+t+\dots+t^{n-1})^d).$$

Note that

$$\mathcal{D}_{i,j} = \text{Card}\{(x_1, \dots, x_d) \in \mathbb{N}^d \mid \forall l, x_l \leq n-1; x_1 + \dots + x_d = in - j\}.$$

Finally let $\mathcal{D}[\sigma, \tau]$ be the infinite square matrix $\mathcal{D}[\sigma, \tau] = (\mathcal{D}_{i+\sigma, j+\tau})$. For $\sigma = \tau = 0$ we write \mathcal{D} instead $\mathcal{D}[0, 0]$. We can give some properties of the numbers $\mathcal{D}_{i,j}$.

Lemma 3.3. *Let $i, j, k \in \mathbb{Z}$, we have:*

- $\mathcal{D}_{i,j} = 0$ if either $in - j < 0$ or $in - j > d(n-1)$.
- For any i, j , $\mathcal{D}_{i,j} = \mathcal{D}_{d-i, d-j}$. That is \mathcal{D} is symmetrical around the point $(d/2, d/2)$.
- For $0 \leq k \leq n-1$, $\mathcal{D}_{d, d+k} = \binom{k+d-1}{d-1}$.
- $\mathcal{D}_{1,0} = \binom{n+d-1}{d-1} - d$, and for $1 \leq k \leq n$, $\mathcal{D}_{1,k} = \binom{n-k+d-1}{d-1}$.
- For any integers q, k , $\mathcal{D}_{d+q, nq+k} = \mathcal{D}_{d,k}$.
- For any i , let $d-i = nq - k$ with $q = \lceil \frac{d-i}{n} \rceil$, $0 \leq k < n$, then

$$\mathcal{D}_{d-\lceil \frac{d-i}{n} \rceil, i} = \binom{k+d-1}{d-1} = \binom{n\lceil \frac{d-i}{n} \rceil + i - 1}{d-1}.$$

Proof. The first claim is trivial. In order to prove the other claims, let remark that the map $(x_1, \dots, x_d) \mapsto (y_1, \dots, y_d)$, where $y_l = (n-1) - x_l$ for $l = 1, \dots, d$, establishes a bijection between

$$\{(x_1, \dots, x_d) \in \mathbb{N}^d \mid x_l \leq n-1 \text{ for } l = 1, \dots, d; \ x_1 + \dots + x_d = in - j\}$$

and

$$\{(y_1, \dots, y_d) \in \mathbb{N}^d \mid y_l \leq n-1 \text{ for } l = 1, \dots, d; \ y_1 + \dots + y_d = (d-i)n - (d-j)\}.$$

The third claim follows from the second claim, because if $0 \leq k \leq n-1$, then the sets

$$\{(x_1, \dots, x_d) \in \mathbb{N}^d \mid x_l \leq n-1 \text{ for } l = 1, \dots, d; \ x_1 + \dots + x_d = dn - d - k\}$$

and

$$\{(y_1, \dots, y_d) \in \mathbb{N}^d \mid y_1 + \dots + y_d = k\}$$

are in bijection.

The fourth claim follows trivially from the previous items.

The fifth claim follows from the equality: $(d+q)n - (nq+k) = dn - k$.

Finally the sixth claim follows from the third claim, since, if $d-i = nq - k$ with $0 \leq k < n$, then $(d-q)n - i = dn - (d+k)$, hence $\mathcal{D}_{d-q,i} = \mathcal{D}_{d,d+k}$, and $n \lceil \frac{d-i}{n} \rceil + i - 1 = k + d - 1$. \square

Remark 3.4. *With the notations introduced in 3.2, it is clear that $f^{<n, kn+\tau>}(t) = t^{-k} f^{<n, \tau>}(t)$, which implies $h^{<n, kn+\tau>}(t) = t^{-k} h^{<n, \tau>}(t)$ for any integer numbers k, τ .*

The following Theorem improves and gives a simpler proof of [2, Theorem 1.1]:

Theorem 3.5. *Fix integers $d, n \in \mathbb{N}^*, \tau \in \mathbb{Z}$. Let $(a_l)_{l \in \mathbb{Z}}$ be a sequence of complex numbers, such that $a_l = 0$ for $l < 0$, set :*

$$f(t) = \sum_{l \in \mathbb{Z}} a_l t^l = \frac{h(t)}{(1-t)^d},$$

$$f^{<n, \tau>}(t) = \sum_{l \in \mathbb{Z}} a_{nl+\tau} t^l = \frac{h^{<n, \tau>}(t)}{(1-t)^d},$$

where $h(t), h^{<n, \tau>}(t) \in \mathbb{C}[t, t^{-1}]$. Then

$$\text{for any } \tau \in \mathbb{Z}, \overrightarrow{h^{<n, kn+\tau>}} = \mathcal{D}[-k, -\tau] \vec{h}.$$

Proof. Because the Remark 3.4 we have to compute $h^{<n, \tau>}(t)$ only for $0 \leq \tau \leq n-1$. The following formula is clear:

$$f^{<n, 0>}(t^n) + t f^{<n, 1>}(t^n) + \dots + t^{n-1} f^{<n, n-1>}(t^n) = f(t),$$

hence

$$\frac{h^{<n, 0>}(t^n) + t h^{<n, 1>}(t^n) + \dots + t^{n-1} h^{<n, n-1>}(t^n)}{(1-t^n)^d} = \frac{h(t)}{(1-t)^d},$$

and

$$h^{<n,0>}(t^n) + th^{<n,1>}(t^n) + \dots + t^{n-1}h^{<n,n-1>}(t^n) = h(t) \frac{(1-t^n)^d}{(1-t)^d},$$

$t^\tau h^{<n,\tau>}(t^n)$ equals the sum of all the terms $A_\beta t^\beta$ of $h(t) \frac{(1-t^n)^d}{(1-t)^d}$ with $\beta \equiv \tau \pmod n$. In particular $h^{<n,\tau>}(t)$ is a linear function of $h(t)$. So it is enough to compute $h^{<n,\tau>}(t)$ for the canonical basis $\{\varepsilon_j := t^j, j \in \mathbb{Z}\}$ of $\mathbb{C}[t, t^{-1}]$. We have

$$[t^i](h^{<n,\tau>}(t)) = [t^{ni+\tau}](h(t) \frac{(1-t^n)^d}{(1-t)^d}),$$

hence

$$\forall j \in \mathbb{Z}; [t^i](\varepsilon_j^{<n,\tau>}(t)) = [t^{ni+\tau}](t^j) \frac{(1-t^n)^d}{(1-t)^d} = [t^{ni+\tau-j}](\frac{(1-t^n)^d}{(1-t)^d})$$

which proves our statement. \square

Corollary 3.6. *Fix an integer $d \in \mathbb{N}^*$. For $j \in \mathbb{Z}$, let $\vec{\varepsilon}_j$ be the h -vector of the polynomial t^j . Then for any $n \in \mathbb{N}^*$, we have $\deg \varepsilon_j^{<n>} = d - \lceil \frac{d-j}{n} \rceil$. Moreover the leading coefficient of $\varepsilon_j^{<n>}$ is $\binom{n \lceil \frac{d-j}{n} \rceil + j - 1}{d-1}$.*

Proof. Let remark that the set of $t^j, j \in \mathbb{Z}$ is the canonical basis of $\mathbb{C}[t, t^{-1}]$. We have by Theorem 3.5 that $\mathcal{D}\vec{\varepsilon}_j = \vec{\varepsilon}_j^{<n>}$, hence $\varepsilon_j^{<n>}$ is the j column vector of \mathcal{D} . By the Example 2.8, we have that $\deg \varepsilon_j^{<n>} = d - \lceil \frac{d-j}{n} \rceil$.

The last claim follows from the Lemma 3.3. Indeed for any $j \in \mathbb{Z}$, we have $\mathcal{D}_{d-\lceil \frac{d-j}{n} \rceil, j} = \binom{n \lceil \frac{d-j}{n} \rceil + j - 1}{d-1}$. This proves that the leading coefficient of $\varepsilon_j^{<n>}$ is $\binom{n \lceil \frac{d-j}{n} \rceil + j - 1}{d-1}$. \square

Example 3.7. *Let $d = 2$ and $n \in \mathbb{N}^*$, we can describe the matrix \mathcal{D}*

ij	$-(n+1)$	\dots	-1	0	1	2	3	\dots	n	$n+1$	$n+2$	\dots	$2n$
-1	2	\dots	0	0	0	0	0	\dots	0	0	0	\dots	0
0	$n-2$	\dots	2	1	0	0	0	\dots	0	0	0	\dots	0
1	0	\dots	$n-2$	$n-1$	n	$n-1$	$n-2$	\dots	1	0	0	\dots	0
2	0	\dots	0	0	0	1	2	\dots	$n-1$	n	$n-1$	\dots	1
3	0	\dots	0	0	0	0	0	\dots	0	0	1	\dots	$n-1$

Theorem 3.8. Fix integers $d, n \in \mathbb{N}^*, \tau \in \mathbb{Z}$. Let $(a_l)_{l \in \mathbb{Z}}$ be a sequence of complex numbers, such that $a_l = 0$ for $l < 0$, set :

$$f(t) = \sum_{l \in \mathbb{Z}} a_l t^l, \quad f^{<n, \tau>}(t) = \sum_{l \in \mathbb{Z}} a_{nl+\tau} t^l.$$

If $f(t) = \frac{h(t)}{(1-t)^d}$ with $h(t) \in \mathbb{C}[t, t^{-1}]$ then $f^{<n, \tau>}(t) = \frac{h^{<n>}(t)}{(1-t)^d}$ with $h^{<n>}(t) \in \mathbb{C}[t, t^{-1}]$ such that:

- $\deg h^{<n, \tau>}(t) \leq d - \lceil \frac{d - \deg h(t) + \tau}{n} \rceil$,
- If all the coefficients of $h(t)$ are positive real numbers then $\deg h^{<n, \tau>}(t) = d - \lceil \frac{d - \deg h(t) + \tau}{n} \rceil$,
- If $\deg h(t) = d$ then $\deg h^{<n>}(t) = d$.

Proof. Let $f(t) = \sum_{l \in \mathbb{Z}} a_l t^l = \frac{h(t)}{(1-t)^d}$, where $h(t) \in \mathbb{C}[t, t^{-1}]$ $h(t) = \gamma_\sigma t^\sigma + \dots + \gamma_s t^s$ with $\deg h(t) = s, \gamma_s \neq 0$. It follows that $\vec{h} = \sum_{l=\sigma}^s \gamma_l \vec{\varepsilon}_l$. We multiply this relation on the left by $\mathcal{D}[-\tau]$, so Theorem 3.5 implies $\vec{h}^{<n, \tau>} = \sum_{l=\sigma}^s \gamma_l \vec{\varepsilon}_{l-\tau}^{<n>}$. Since $\deg \vec{\varepsilon}_{\sigma-\tau}^{<n>} \leq \deg \vec{\varepsilon}_{\sigma-\tau+1}^{<n>} \leq \dots \leq \deg \vec{\varepsilon}_{s-\tau}^{<n>}$, we have, $\deg \vec{h}^{<n, \tau>} \leq \deg \vec{\varepsilon}_{s-\tau}^{<n>}$. It is clear that if all the coefficients of $h(t)$ are positive real numbers then $\deg \vec{h}^{<n, \tau>} = \deg \vec{\varepsilon}_{s-\tau}^{<n>}$.

In the special case $s = d$, we have seen that for $0 \leq l \leq d-1$ and any $n \in \mathbb{N}^*$, $\deg \vec{\varepsilon}_l^{<n>} = d - \lceil \frac{d-l}{n} \rceil \leq d-1$, and $\deg \vec{\varepsilon}_d^{<n>} = d$, which implies $\deg \vec{h}^{<n>} = d$. \square

Theorem 3.9. Let $n \in \mathbb{N}^*$, S be a standard graded polynomial ring, $M = \oplus_{l \in \mathbb{Z}} M_l$ be a finitely generated Cohen-Macaulay graded S -module of dimension $d \geq 1$, and $M^{<n>} = \oplus_{l \in \mathbb{Z}} M_{nl}$. Let $\frac{Q(t)}{(1-t)^d}$ be the Hilbert-Poincaré series of M , where $Q(t) = \gamma_\sigma t^\sigma + \dots + \gamma_s t^s \in \mathbb{C}[t, t^{-1}]$ is the h -polynomial of M , with $\text{reg}(M) = \deg Q(t) = s$. Then

- $\text{reg } M^{<n>} = d - \lceil \frac{d - \text{reg } M}{n} \rceil$. Moreover by taking the sum over all index l such that $\lceil \frac{d-l}{n} \rceil = \lceil \frac{d - \text{reg } M}{n} \rceil$, we will get the leading coefficient of $Q^{<n>}(t)$:

$$\sum_{l \mid \lceil \frac{d-l}{n} \rceil = \lceil \frac{d - \text{reg } M}{n} \rceil} \gamma_l \binom{n \lceil \frac{d-l}{n} \rceil + l - 1}{d-1}.$$

- If $\text{reg } M \leq d - 1$ and $n \geq d$ then $\text{reg } M^{<n>} = d - 1$, and the leading coefficient of $Q^{<n>}(t)$ is

$$\sum_{l=0}^{d-1} \gamma_l \binom{n \lceil \frac{d-l}{n} \rceil + l - 1}{d-1}.$$

- If $n > \text{reg } M \geq d$ then $\text{reg } M^{<n>} = d$, and the leading coefficient of $Q^{<n>}(t)$ is

$$\sum_{l=d}^{\text{reg } M} \gamma_l \binom{l-1}{d-1}.$$

Proof. We have $\vec{Q} = \sum_{l=\sigma}^s \gamma_l \vec{\varepsilon}_l$. We multiply this relation on the left by \mathcal{D} , so Theorem 3.5 implies that for any $n \in \mathbb{N}^*$, $\overrightarrow{Q^{<n>}} = \sum_{l=\sigma}^s \gamma_l \overrightarrow{\varepsilon_l^{<n>}}$. Since $\gamma_l \geq 0$ for all l , $\gamma_s > 0$, and $\deg \overrightarrow{\varepsilon_\sigma^{<n>}} \leq \deg \overrightarrow{\varepsilon_{\sigma+1}^{<n>}} \leq \dots \leq \deg \overrightarrow{\varepsilon_s^{<n>}}$, we have, $\deg \overrightarrow{Q^{<n>}} = \deg \overrightarrow{\varepsilon_s^{<n>}} = d - \lceil \frac{d - \text{reg } M}{n} \rceil$, this number is $\text{reg}(M^{<n>})$ since $M^{<n>}$ is a Cohen-Macaulay S -module. The computation of the leading coefficient of $Q^{<n>}(t)$ is immediate from Lemma 3.6. \square

4 h -vector of the Segre-Veronese embedding.

The next Theorem improves partially [1] and [5].

Theorem 4.1. *Let consider the Segre-Veronese ring $R_{\underline{b}, \underline{n}}$, $\dim R_{\underline{b}, \underline{n}} = b_1 + \dots + b_m + 1$. Let $P_{R_{\underline{b}, \underline{n}}}(t) = \frac{Q_{R_{\underline{b}, \underline{n}}}(t)}{(1-t)^{\dim R_{\underline{b}, \underline{n}}}}$ be the Hilbert-Poincaré series of $R_{\underline{b}, \underline{n}}$, with $Q_{R_{\underline{b}, \underline{n}}}(t) = h_0 + h_1 t + \dots + h_{r_{\underline{b}, \underline{n}}} t^{r_{\underline{b}, \underline{n}}}$, where $r_{\underline{b}, \underline{n}} = \deg Q_{R_{\underline{b}, \underline{n}}}(t)$ is the Castelnuovo-Mumford regularity of $R_{\underline{b}, \underline{n}}$. We set $\alpha_{\underline{b}, \underline{n}} = \dim R_{\underline{b}, \underline{n}} - r_{\underline{b}, \underline{n}}$. After a permutation of b_1, \dots, b_m , we can assume that $\lceil \frac{b_1+1}{n_1} \rceil > \frac{b_i}{n_i} \forall i$, then*

$$\alpha_{\underline{b}, \underline{n}} = \lceil \frac{b_1+1}{n_1} \rceil \quad r_{\underline{b}, \underline{n}} = (b_1 + \dots + b_m + 1) - \lceil \frac{b_1+1}{n_1} \rceil,$$

and the highest Betti number of $R_{\underline{b}, \underline{n}}$ is

$$\beta_{r_{\underline{b}, \underline{n}}} = h_{r_{\underline{b}, \underline{n}}} = \binom{n_1 \alpha_{\underline{b}, \underline{n}} - 1}{b_1} \dots \binom{n_m \alpha_{\underline{b}, \underline{n}} - 1}{b_m}$$

Proof. The proof is by double induction on m and b_m . The case $m = 1$ is given by the Example 2.8 and Corollary 3.6, so we can assume $m \geq 2$. We have that $\lceil \frac{b_1+1}{n_1} \rceil > \frac{b_m}{n_m} > \frac{b_m-1}{n_m}$, so by induction hypothesis the theorem is true for $R_{\underline{b}-\epsilon_m, \underline{n}}$, where $\underline{b}-\epsilon_m = (b_1, \dots, b_{m-1}, b_m-1)$. On the other hand the Hilbert function of $R_{\underline{b}, \underline{n}}$ is $H_{R_{\underline{b}, \underline{n}}}(l) = \binom{n_1 l + b_1}{b_1} \dots \binom{n_m l + b_m}{b_m}$, so

$$H_{R_{\underline{b}, \underline{n}}}(l) = (1 + \frac{n_m}{b_m} l) H_{R_{\underline{b}-\epsilon_m, \underline{n}}}(l). \quad (1)$$

Let $P_{R_{\underline{b}-\epsilon_m, \underline{n}}}(t) = \frac{Q_{R_{\underline{b}-\epsilon_m, \underline{n}}}(t)}{(1-t)^{b_1+\dots+b_m}}$ be the Hilbert-Poincaré series of $R_{\underline{b}-\epsilon_m, \underline{n}}$, where $Q_{R_{\underline{b}-\epsilon_m, \underline{n}}}(t) = h_0 + h_1 t + \dots + h_{r_{\underline{b}-\epsilon_m, \underline{n}}} t^{r_{\underline{b}-\epsilon_m, \underline{n}}}$, with $h_{r_{\underline{b}-\epsilon_m, \underline{n}}} \neq 0$. In order to avoid any confusion we also set : $P_{R_{\underline{b}, \underline{n}}}(t) = \frac{Q_{R_{\underline{b}, \underline{n}}}(t)}{(1-t)^{b_1+\dots+b_m+1}}$ be the Hilbert-Poincaré series of $R_{\underline{b}, \underline{n}}$, where

$$Q_{R_{\underline{b}, \underline{n}}}(t) = \hat{h}_0 + \dots + \hat{h}_{r_{\underline{b}, \underline{n}}} t^{r_{\underline{b}, \underline{n}}},$$

with $\hat{h}_{r_{\underline{b}, \underline{n}}} \neq 0$.

Let $\beta = \frac{n_m}{b_m}$, by simple calculations from (1) we get:

$$P_{R_{\underline{b}, \underline{n}}}(t) = P_{R_{\underline{b}-\epsilon_m, \underline{n}}}(t) + \beta t P'_{R_{\underline{b}-\epsilon_m, \underline{n}}}(t). \quad (2)$$

Hence $\dim R_{\underline{b}, \underline{n}} = \dim R_{\underline{b}-\epsilon_m, \underline{n}} + 1$, and

$$Q_{R_{\underline{b}, \underline{n}}}(t) = Q_{R_{\underline{b}-\epsilon_m, \underline{n}}}(t) + t[Q_{R_{\underline{b}-\epsilon_m, \underline{n}}}(t)(\beta R_{\underline{b}-\epsilon_m, \underline{n}} - 1) + \beta Q'_{R_{\underline{b}-\epsilon_m, \underline{n}}}(t) - \beta t Q'_{R_{\underline{b}-\epsilon_m, \underline{n}}}(t)],$$

note that $Q_{R_{\underline{b}, \underline{n}}}(1) = \beta \dim R_{\underline{b}-\epsilon_m, \underline{n}} Q_{R_{\underline{b}-\epsilon_m, \underline{n}}}(1) \neq 0$.

In particular we have $r_{\underline{b}, \underline{n}} \leq r_{\underline{b}-\epsilon_m, \underline{n}} + 1$ and for all $k = 0, \dots, r_{\underline{b}-\epsilon_m, \underline{n}} + 1$ we have

$$\hat{h}_k = h_{k-1} \left(\frac{n_m}{b_m} \dim R_{\underline{b}-\epsilon_m, \underline{n}} - (k-1) \frac{n_m}{b_m} - 1 \right) + h_k \left(k \frac{n_m}{b_m} + 1 \right). \quad (3)$$

By induction hypothesis we have $\alpha_{\underline{b}-\epsilon_m, \underline{n}} = \lceil \frac{b_1+1}{n_1} \rceil \neq \frac{b_m}{n_m}$, so we put $k = r_{\underline{b}-\epsilon_m, \underline{n}} + 1$ in equality (3), and we get:

$$\hat{h}_{r_{\underline{b}-\epsilon_m, \underline{n}}+1} = h_{r_{\underline{b}-\epsilon_m, \underline{n}}} \left(\frac{n_m \alpha_{\underline{b}-\epsilon_m, \underline{n}} - b_m}{b_m} \right) \neq 0.$$

Hence $\hat{h}_{r_{\underline{b}-\epsilon_m, \underline{n}}+1}$ is the leading coefficient of $Q_{R_{\underline{b}, \underline{n}}}$ and $r_{\underline{b}, \underline{n}} = r_{\underline{b}-\epsilon_m, \underline{n}} + 1$ and $\alpha_{\underline{b}, \underline{n}} = \alpha_{\underline{b}-\epsilon_m, \underline{n}} = \lceil \frac{b_1+1}{n_1} \rceil$. By induction hypothesis

$$h_{r_{\underline{b}-\epsilon_m, \underline{n}}} = \binom{n_1 \alpha_{\underline{b}, \underline{n}} - 1}{b_1} \dots \binom{n_{m-1} \alpha_{\underline{b}, \underline{n}} - 1}{b_{m-1}} \binom{n_m \alpha_{\underline{b}, \underline{n}} - 1}{b_m}$$

so that

$$\begin{aligned}\hat{h}_{r_{\underline{b}, \underline{n}}} &= \binom{n_1 \alpha_{\underline{b}, \underline{n}} - 1}{b_1} \cdots \binom{n_{m-1} \alpha_{\underline{b}, \underline{n}} - 1}{b_{m-1}} \binom{n_m \alpha_{\underline{b}, \underline{n}} - 1}{b_m - 1} \left(\frac{n_m \alpha_{\underline{b}, \underline{n}} - b_m}{b_m} \right) \\ &= \binom{n_1 \alpha_{\underline{b}, \underline{n}} - 1}{b_1} \cdots \binom{n_{m-1} \alpha_{\underline{b}, \underline{n}} - 1}{b_{m-1}} \binom{n_m \alpha_{\underline{b}, \underline{n}} - 1}{b_m}.\end{aligned}$$

□

5 Rings of Veronese type

Let $b, n \in \mathbb{N}^*$, $\underline{a} = (a_0, \dots, a_b) \in \mathbb{N}^{b+1}$ such that $1 \leq a_i \leq n$, $a_0 + \dots + a_b > n$, and $\mathcal{M}_{b, n, \underline{a}}$ be the set of monomials of the polynomial ring $K[x_0, \dots, x_b]$:

$$\mathcal{M}_{b, n, \underline{a}} = \{x_0^{\alpha_0} \dots x_b^{\alpha_b} \mid \alpha_0 + \dots + \alpha_b = n, \alpha_i \leq a_i, \forall i = 0, \dots, b\}.$$

Let denote by $R_{b, n, \underline{a}}$ the toric subring of $K[x_0, \dots, x_b]$ generated by $\mathcal{M}_{b, n, \underline{a}}$. It is well known that $R_{b, n, \underline{a}}$ is a Cohen-Macaulay ring. Let \mathcal{S} be the collection of subsets of $\{0, \dots, b\}$ such that : $S \in \mathcal{S}$ if and only if $S \subset \{0, \dots, b\}$, and $\Sigma S := \sum_{i \in S} a_i < n$.

Theorem 5.1. ([9]) *With the above notations the Hilbert-function of $R_{b, n, \underline{a}}$ is*

$$\forall l \geq 0; \quad H_{b, n, \underline{a}}(l) = \sum_{S \in \mathcal{S}} (-1)^{|S|} \binom{l(n - \Sigma S) - |S| + b}{b}$$

We have $\dim(R_{b, n, \underline{a}}) = b + 1$, and its degree or multiplicity is

$$\deg(R_{b, n, \underline{a}}) = \sum_{S \in \mathcal{S}} (-1)^{|S|} (n - \Sigma S)^b.$$

Our aim is to study the Hilbert-Poincaré series of $R_{b, n, \underline{a}}$:

$$P_{R_{b, n, \underline{a}}} = \sum_{S \in \mathcal{S}} (-1)^{|S|} \sum_{l \geq 0} \binom{l(n - \Sigma S) - |S| + b}{b} t^l.$$

The following Corollary follows immediately from 3.6.

Corollary 5.2. *For any $S \in \mathcal{S}$ and $k \in \mathbb{N}^*$, we have:*

$$\sum_{l \geq 0} \binom{kl - |S| + b}{b} t^l = \frac{Q_{S, k}(t)}{(1 - t)^{b+1}},$$

where $Q_{S,k}(t)$ is a polynomial with $Q_{S,k}(1) \neq 0$, with leading term

$$\binom{k\alpha_{S,k} + |S| - 1}{b} t^{b+1-\alpha_{S,k}},$$

with $\alpha_{S,k} = \lceil \frac{b+1-|S|}{k} \rceil$.

The following theorem is immediate from 5.1 and Corollary 5.2. It improves the description of the Hilbert Poincaré series given in [9].

Theorem 5.3. *With the above notations, let \mathcal{S} be the collection of subsets of $\{0, \dots, b\}$ such that : $S \in \mathcal{S}$ if and only if $S \subset \{0, \dots, b\}$, and $\Sigma S := \sum_{i \in S} a_i < n$. Then we can write the Hilbert-Poincaré series of $R_{b,n,\underline{a}}$:*

$$P_{R_{b,n,\underline{a}}} = \frac{Q_{b,n,\underline{a}}(t)}{(1-t)^{b+1}},$$

with $Q_{b,n,\underline{a}}(t) = \sum_{S \in \mathcal{S}} (-1)^{|S|} Q_{S,n}(t)$, where $Q_{S,n}(t)$ is a polynomial with $Q_{S,n}(1) \neq 0$, with leading term

$$\binom{(n - \Sigma S)\alpha_{S,n-\Sigma S} + |S| - 1}{b} t^{b+1-\alpha_{S,n-\Sigma S}},$$

where $\alpha_{S,n-\Sigma S} = \lceil \frac{b+1-|S|}{n-\Sigma S} \rceil$.

Part one of the following corollary improves [9][Cor. 2.12].

Corollary 5.4. *With the above notations:*

1. $\text{reg}(R_{b,n,\underline{a}}) \leq b + 1 - \lceil \frac{b+1}{n} \rceil$, and the equality is true if and only if

$$\sum_{S \in \mathcal{S}, \alpha_{S,n-\Sigma S} = \lceil \frac{b+1}{n} \rceil} (-1)^{|S|} \binom{(n - \Sigma S)\alpha_{S,n-\Sigma S} + |S| - 1}{b} \neq 0$$

2. If $b + 1 > n^2$ then $\text{reg}(R_{b,n,\underline{a}}) = b + 1 - \lceil \frac{b+1}{n} \rceil$. Moreover the leading term of $Q_{b,n,\underline{a}}(t)$ is $\binom{(n \lceil \frac{b+1}{n} \rceil - 1)}{b} t^{b+1-\lceil \frac{b+1}{n} \rceil}$,

Proof. 1. It is enough to prove that $\min_{S \in \mathcal{S}} \lceil \frac{b+1-|S|}{n-\Sigma S} \rceil = \lceil \frac{b+1}{n} \rceil$. We consider two cases,

- if $b + 1 < n$ then $\lceil \frac{b+1}{n} \rceil = 1 \leq \lceil \frac{b+1-|S|}{n-\Sigma S} \rceil, \forall S \in \mathcal{S}$.

- If $b + 1 \geq n$, then

$$\frac{b+1}{n} \leq \frac{b+1-|S|}{n-\Sigma S} \Leftrightarrow (b+1)(n-\Sigma S) \leq n(b+1-|S|) \Leftrightarrow (b+1)\Sigma S \geq n|S|,$$

this is true since by hypothesis $\frac{b+1}{n} \geq 1 \geq \frac{|S|}{\Sigma S}$.

2. Let $b + 1 > n^2$ and $S \neq \emptyset$. By definition $\lceil \frac{b+1}{n} \rceil$ is the integer q such that $b + 1 = qn - r$, with $0 \leq r < n$ and $q \geq n + 1$. We have

$$b + 1 - |S| = qn - r - |S| = q(n - \Sigma S) - r - |S| + q\Sigma S,$$

and $q\Sigma S - |S| \geq (n + 1)\Sigma S - |S| \geq n\Sigma S > r$, so that $q\Sigma S - |S| - r > 0$, hence $\lceil \frac{b+1-|S|}{n-\Sigma S} \rceil > q = \lceil \frac{b+1}{n} \rceil$. \square

In general leading terms of the alternating sum can cancel, as we can see in the next example.

Example 5.5. Let consider the ring $R_{4,3,(1,1,1,1,1)}$, the sets S can have 0, 1 or 2 elements, and we have: If $S = \emptyset$ then $\alpha_{\emptyset,3} = \lceil \frac{5}{3} \rceil = 2$, if $|S| = 1$ then $\alpha_{S,3} = \lceil \frac{4}{2} \rceil = 2$, and finally if $|S| = 2$ then $\alpha_{S,2} = \lceil \frac{3}{1} \rceil = 3$. By using Theorem 5.3 we can write

$$P_{R_{4,3,(1,1,1,1,1)}} = \frac{Q_0(t) - 5Q_1(t) + 10Q_2(t)}{(1-t)^5},$$

, with $Q_0(t) = 5t^3 + \dots$; $Q_1(t) = t^3 + \dots$; $Q_2(t) = t^2 + \dots$. Note that in this case $Q_0(t) - 5Q_1(t) + 10Q_2(t) = h_0 + h_1t + h_2t^2$, where $h_0 = 1, h_1 = 5$ and since $h_0 + h_1 + h_2 = \deg(R_{4,3,(1,1,1,1,1)}) = 11$, we get $h_2 = 5$, so that

$$P_{R_{4,3,(1,1,1,1,1)}} = \frac{1 + 5t + 5t^2}{(1-t)^5}.$$

References

- [1] Barcanescu, S.; Manolache, N., Betti numbers of Segre-Veronese singularities. Rev. Roum. Math. Pures Appl. 26, 549-565 (1981).

- [2] Breni, F. ; Welker V., The Veronese construction for formal power series and graded algebras *Adv. Applied Math.*, 42 (2009), 545-556.
- [3] Brodmann, M.P.; Sharp, R.Y., Local cohomology. An algebraic introduction with geometric applications. Cambridge Studies in Advanced Mathematics. 60. Cambridge: Cambridge University Press. xv, 416 p. (1998).
- [4] Bruns, Winfried; Herzog, Jürgen, Cohen-Macaulay rings. Rev. ed. Cambridge Studies in Advanced Mathematics. 39. Cambridge: Cambridge University Press. xiv, 453 p. (1998).
- [5] Cox, David A.; Materov, Evgeny, Regularity and Segre-Veronese embeddings. *Proc. Am. Math. Soc.* 137, No. 6, 1883-1890 (2009).
- [6] De Negri, Emanuela; Hibi, Takayuki, Gorenstein algebras of Veronese type. *J. Algebra* 193, No.2, 629-639 (1997).
- [7] Eisenbud, David, The geometry of syzygies. A second course in commutative algebra and algebraic geometry. Graduate Texts in Mathematics 229. New York, NY: Springer. xvi, 243 p. (2005).
- [8] Goto, Shiro and Watanabe, Keiichi, On graded rings, I. *J. Math. Soc. Japan* Volume 30, Number 2 (1978), 179-213.
- [9] Katzman, Mordechai, The Hilbert series of algebras of the Veronese type. *Commun. Algebra* 33, No. 4, 1141-1146 (2005).
- [10] Morales M., *Fonctions de Hilbert, genre géométrique d'une singularité quasi-homogène Cohen-Macaulay*. CRAS Paris, t.301, série A n° 14 (1985).